

For the use of Registered Medical Practitioner or Hospital use or Laboratory use only

Bricanyl® Syrup

Composition

Each ml of syrup contains:
Terbutaline Sulphate I.P. 0.3 mg.

Description: Terbutaline (Bricanyl Syrup) is an adrenergic agonist which selectively stimulates beta-2-receptors, thus producing relaxation of bronchial smooth muscle, inhibition of the release of endogenous spasmogens, inhibition of oedema caused by endogenous mediators and increased mucociliary clearance.

Indications: For the relief of bronchospasm in bronchial asthma, chronic bronchitis, emphysema and other diseases of the lung where bronchoconstriction is a complicating factor.

Precautions: Patients with thyrotoxicosis should use terbutaline with caution. Terbutaline should be given with care-as any other sympathomimetic amine - to patients with cardiovascular diseases or diabetes and to patients with a record of hypersensitivity to sympathomimetic amines.

Adverse Effects: The frequency of side effects with oral terbutaline is low. Those recorded e.g. tremor and palpitations are characteristic of all sympathomimetic amines. Tremor diminishes on continued treatment. Tremor and palpitations subside if the dose is lowered.

Dosage: Dosage should be individualized

Adults : Bricanyl® Syrup (0.3mg/ml), 10-15 ml, 2-3 times a day

Children : 0.075mg (0.25 ml)/kg. body weight), 3 times in a 24 hour period.

The following serves as a guideline.

7-15 years : Bricanyl® Syrup, 5-10ml 2-3 times a day (1-2 teaspoonful).

3-6 years : Bricanyl® Syrup, 2.5-5 ml 2-4 times a day (1/2-1 teaspoonful).

2 years : Bricanyl® Syrup, 2.5 ml 2-3 times a day (1/2 teaspoonful)

Presentation : Bricanyl® Syrup (0.3mg/ml), 60 ml bottle

For further information please contact:

AstraZeneca Pharma India Ltd.
Bellary Road, Hebbal
Bangalore – 560 024